

On the zeros of the Hahn-Exton q -Bessel function and associated q -Lommel polynomials

H.T. Koelink*

R.F. Swarttouw†

1 Introduction

For the Bessel function

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{z}{2}\right)^{\nu+2k}}{k! \Gamma(\nu+1+k)} \quad (1.1)$$

there exist several q -analogues. The oldest q -analogues of the Bessel function were introduced by F.H. Jackson at the beginning of this century, see M.E.H. Ismail [4] for the appropriate references. Another q -analogue of the Bessel function has been introduced by W. Hahn in a special case and by H. Exton in full generality, see R.F. Swarttouw [9] for a historic overview.

Here we concentrate on properties of the Hahn-Exton q -Bessel function and in particular on its zeros and the associated q -Lommel polynomials. Ismail has proved very satisfactory results on this subject for the Jackson q -Bessel functions, cf. [4]. In particular, he proves orthogonality relations for the associated q -Lommel polynomials.

In section 2 we present the definition of the Hahn-Exton q -Bessel function and some of its properties. The zeros of the Hahn-Exton q -Bessel function of order $\nu > -1$ are the subject of section 3. The proofs of the statements in this section rest on the evaluation of a q -integral closely related to the Fourier-Bessel orthogonality relations for the Hahn-Exton q -Bessel function. The results of section 3 are in accordance with the results on the zeros of the Bessel function. Section 4 deals with two kinds of associated q -Lommel polynomials for which we present explicit forms as well as some properties. However, the first type does not give rise to orthogonal polynomials, while the second type provides polynomials, which are closely related to the orthogonal modified q -Lommel polynomials found by M.E.H. Ismail.

Comparison of our results with the results for the Jackson q -Bessel functions shows that the Hahn-Exton q -Bessel function is less similar to the Bessel function in this respect than the Jackson q -Bessel function, since we are not able to prove new orthogonality relations for the q -Lommel polynomials associated with the Hahn-Exton q -Bessel function. This seems in support of M. Rahman [7], who favours the Jackson q -Bessel function over the

*Van Swietenstraat 3, 2334 EA Leiden, The Netherlands

†Vrije Universiteit, Faculteit Wiskunde en Informatica, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands

Hahn-Exton q -Bessel function. However, it should be noted that the proofs concerning the zeros of the Hahn-Exton q -Bessel function are much simpler and more direct than those for the Jackson q -Bessel function.

In our opinion both q -analogues of the Bessel function are interesting functions and possess nice properties. Moreover, from the harmonic analysis on the quantum group of plane motions, cf. H.T. Koelink [5, §6.7], it follows that there exists a common generalisation of the Jackson and Hahn-Exton q -Bessel function.

2 The Hahn-Exton q -Bessel function

In this section we present the Hahn-Exton q -Bessel function and some of its properties. References for these results are T.H. Koornwinder and R.F. Swarttouw [6] and R.F. Swarttouw [9]. The q -Bessel functions are defined in terms of basic hypergeometric series, which we will briefly recall. More information on basic hypergeometric series can be found in the book by G. Gasper and M. Rahman [3, Chapter 1].

We fix $0 < q < 1$ and we define the q -shifted factorials

$$(a; q)_k = \begin{cases} 1 & \text{if } k = 0 \\ (1-a)(1-aq) \cdots (1-aq^{k-1}) & \text{if } k \geq 1 \end{cases} \quad (2.1)$$

for arbitrary $a \in \mathbb{C}$ and $k \in \mathbb{Z}_+$. The limit $k \rightarrow \infty$ is well defined and yields

$$(a; q)_\infty = \lim_{k \rightarrow \infty} (a; q)_k. \quad (2.2)$$

The q -hypergeometric series (or basic hypergeometric) series ${}_r\phi_s$ is defined by

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q; z \right) = \sum_{k=0}^{\infty} \frac{(a_1; q)_k (a_2; q)_k \cdots (a_r; q)_k}{(q; q)_k (b_1; q)_k \cdots (b_s; q)_k} \left((-1)^k q^{\frac{1}{2}k(k-1)} \right)^{1+s-r} z^k \quad (2.3)$$

whenever the series converges.

The Hahn-Exton q -Bessel function $J_\nu(x; q)$ of order ν is defined as

$$J_\nu(x; q) = x^\nu \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} {}_1\phi_1 \left(\begin{matrix} 0 \\ q^{\nu+1} \end{matrix} \middle| q; qx^2 \right). \quad (2.4)$$

From (2.3) we see that $x^{-\nu} J_\nu(x; q^2)$ defines a non-zero analytic function on $\mathbb{C}$, so that $J_\nu(x; q)$ is analytic in $\mathbb{C} \setminus \{0\}$. The Hahn-Exton q -Bessel function is a q -analogue of the Bessel function, since $J_\nu((1-q)x; q)$ tends to the Bessel function $J_\nu(2x)$ of order ν as $q \uparrow 1$, cf. [9, Appendix A]. The standard text on Bessel functions is the classic by G.N. Watson [11].

Next we introduce two concepts of q -analysis: the q -derivative and the q -integral, cf. [3, Chapter 1]. The q -derivative of a function f is defined by

$$(D_q f)(x) = \frac{f(x) - f(qx)}{(1-q)x}, \quad x \neq 0. \quad (2.5)$$

Note that the $(D_q f)(x)$ tends to $f'(x)$ for $q \uparrow 1$ whenever f is differentiable at x . The product rule for the q -derivative is

$$(D_q f g)(x) = f(x) (D_q g)(x) + g(qx) (D_q f)(x). \quad (2.6)$$

The q -integral of a function f is defined by

$$\int_0^z f(x) d_q x = (1-q)z \sum_{k=0}^{\infty} f(zq^k) q^k, \quad z > 0. \quad (2.7)$$

For a Riemann-integrable function f this expression tends to $\int_0^z f(x) dx$ as $q \uparrow 1$. It is easily checked that for continuous f

$$\int_0^z (D_q f)(x) d_q x = f(z) - f(0). \quad (2.8)$$

The q -product rule leads to the following q -partial integration rule

$$\int_0^z f(qx) (D_q g)(x) d_q x = f(z)g(z) - f(0)g(0) - \int_0^z (D_q f)(x)g(x) d_q x. \quad (2.9)$$

We will use the following q -derivatives, cf. [9, (3.2.22), (3.2.19)]:

$$D_q \left[(\cdot)^\nu J_\nu(\cdot; q^2) \right] (x) = \frac{x^\nu}{1-q} J_{\nu-1}(x; q^2), \quad (2.10)$$

$$D_q \left[(\cdot)^{-\nu} J_\nu(\cdot; q^2) \right] (x) = -\frac{q^{1-\nu} x^{-\nu}}{1-q} J_{\nu+1}(xq; q^2). \quad (2.11)$$

These formulas are q -analogues of the case $m = 1$ of [11, 3.2(5), 3.2(6)].

The second order differential equation for the Bessel function, cf. [11, 3.2(1)], has the following second order q -difference equation as a q -analogue for the Hahn-Exton q -Bessel function, cf. [9, (4.3.1)]:

$$J_\nu(xq^2; q^2) + q^{-\nu} (x^2 q^2 - 1 - q^{2\nu}) J_\nu(xq; q^2) + J_\nu(x; q^2) = 0. \quad (2.12)$$

It will also be useful to have the following relations at hand. They involve Hahn-Exton q -Bessel functions of several orders and can be found in [9, (3.2.15), (3.2.18), (3.2.21)]:

$$J_{\nu+1}(x; q^2) = \left(\frac{1-q^{2\nu}}{x} + x \right) J_\nu(x; q^2) - J_{\nu-1}(x; q^2), \quad (2.13)$$

$$J_{\nu+1}(xq; q^2) = q^{-\nu-1} \left(\frac{1-q^{2\nu}}{x} J_\nu(x; q^2) - J_{\nu-1}(x; q^2) \right). \quad (2.14)$$

3 On the zeros of the Hahn-Exton q -Bessel function

In this section we prove some results on the zeros of the Hahn-Exton q -Bessel function. The proofs rest on an explicit evaluation of a q -integral related to the Fourier-Bessel orthogonality relations for the Hahn-Exton q -Bessel function and on some formulas relating Hahn-Exton q -Bessel functions of different order. At the end of this section we also present a result on the zeros of the q -derivative of the Hahn-Exton q -Bessel function.

The starting point in the derivation of the results on the zeros of the Hahn-Exton q -Bessel function is the following proposition. It is closely related to the Fourier-Bessel orthogonality relations for the Hahn-Exton q -Bessel functions, originally proved by H. Exton (see [1] and [2]) using Sturmian methods.

Proposition 3.1. *For $\Re(\nu) > -1$, $z > 0$ and $a, b \in \mathbb{C} \setminus \{0\}$ we have*

$$\begin{aligned} & (a^2 - b^2) \int_0^z x J_\nu(ax; q^2) J_\nu(bx; q^2) d_q x \\ &= (1 - q) q^{\nu-1} z \left(a J_{\nu+1}(az; q^2) J_\nu(bz; q^2) - b J_{\nu+1}(bz; q^2) J_\nu(az; q^2) \right). \end{aligned}$$

Proof: Use the q -partial integration rule (2.9) and the formulas (2.10) and (2.11) to obtain

$$\begin{aligned} & \int_0^z x J_\nu(ax; q^2) J_\nu(bx; q^2) d_q x \\ &= \frac{(1 - q)}{b} q^{\nu-1} z J_\nu(az; q^2) J_{\nu+1}(bz; q^2) + \frac{a}{b} \int_0^z x J_{\nu+1}(ax; q^2) J_{\nu+1}(bx; q^2) d_q x. \end{aligned} \tag{3.1}$$

The restriction $\Re(\nu) > -1$ is necessary to ensure the absolute convergence of the series defining the q -integral, cf. (2.7), since the Hahn-Exton q -Bessel function $J_\nu(x; q^2)$ behaves like x^ν times a constant for small x . Interchanging a and b in (3.1) yields a set of two equations, which can be solved easily. $\square$

The proof of Proposition 3.1 is also well known for the Bessel function, see e.g. I.N. Sneddon [8, Chapter 2].

Corollary 3.2. *The zeros of $J_\nu(\cdot; q^2)$, with $\nu > -1$, are real.*

Proof: Suppose $a \neq 0$ is a zero of $J_\nu(\cdot; q^2)$. Since ν is real we have

$$J_\nu(\bar{a}; q^2) = \overline{J_\nu(a; q^2)} = 0.$$

Proposition 3.1 with $z = 1$ and $b = \bar{a}$ yields

$$(a^2 - \bar{a}^2) \int_0^1 x \left| J_\nu(ax; q^2) \right|^2 d_q x = 0. \tag{3.2}$$

Now $a^2 = \bar{a}^2$ if and only if $a \in \mathbb{R}$ or $a \in i\mathbb{R}$, so that in all other cases the q -integral in (3.2) is zero. Using the definition of the q -integral (2.7) we get

$$J_\nu(aq^{k+1}; q^2) = 0, \quad k \in \mathbb{Z}_+,$$

and this implies that $x^{-\nu} J_\nu(x; q^2)$ is identically zero, since it defines an analytic function on $\mathbb{C}$. Hence, $J_\nu(\cdot; q^2) \equiv 0$.

Finally, we have to show that ia , with $a \in \mathbb{R}$, is not a zero of $J_\nu(\cdot; q^2)$. Now by (2.3) and (2.4)

$$J_\nu(ia; q^2) = (ia)^\nu \frac{(q^{2\nu+2}; q^2)_\infty}{(q^2; q^2)_\infty} \sum_{k=0}^{\infty} \frac{q^{k(k+1)} a^{2k}}{(q^2; q^2)_k (q^{2\nu+2}; q^2)_k}$$

and for $\nu > -1$ this expression cannot be zero, since every term in the sum is positive. $\square$

From the series representation (2.4) for the Hahn-Exton q -Bessel function it follows that if a is a zero of $J_\nu(\cdot; q^2)$, then $-a$ is also a zero of $J_\nu(\cdot; q^2)$. Hence we will restrict ourselves to the positive zeros of the Hahn-Exton q -Bessel function of order $\nu > -1$.

To obtain an expression for the q -integral in Proposition 3.1 with $a = b$, we use l'Hôpital's rule. The result is

$$\begin{aligned} \int_0^z x \left(J_\nu(axq; q^2) \right)^2 d_q x &= \frac{(1-q)q^{\nu-1}z}{-2a} \left(az J_{\nu+1}(aqz; q^2) J'_\nu(az; q^2) \right. \\ &\quad \left. - J_{\nu+1}(aqz; q^2) J_\nu(az; q^2) - aqz J'_{\nu+1}(aqz; q^2) J_\nu(az; q^2) \right). \end{aligned}$$

This formula simplifies to

$$\int_0^1 x \left(J_\nu(axq; q^2) \right)^2 d_q x = -\frac{1}{2}(1-q)q^{\nu-1} J_{\nu+1}(aq; q^2) J'_\nu(a; q^2) \quad (3.3)$$

for $z = 1$ and $a \neq 0$ a (real) zero of $J_\nu(\cdot; q^2)$.

Lemma 3.3. *The non-zero (real) zeros of $J_\nu(\cdot; q^2)$, with $\nu > -1$, are simple zeros.*

Proof: Let a be a non-zero (real) zero of $J_\nu(\cdot; q^2)$, with $\nu > -1$. The integral

$$\int_0^1 x \left| J_\nu(axq; q^2) \right|^2 d_q x = \int_0^1 x \left(J_\nu(axq; q^2) \right)^2 d_q x$$

is strictly positive. (If it were zero, this would imply that the Hahn-Exton q -Bessel function is identically zero as in the proof of Corollary 3.2.) Hence, (3.3) implies that $J'_\nu(a; q^2) \neq 0$, which proves the lemma. $\square$

In the proof of this lemma we explicitly use the (usual) derivative of the Hahn-Exton q -Bessel function in (3.3). It shows that the intermingling of (ordinary) analysis and q -analysis may be fruitful and should not be opposed to as M. Rahman [7] proposes.

We now come to one of the main results of this section, which shows that the zeros of the Hahn-Exton q -Bessel function behave like the zeros of the classical Bessel function, cf. Watson [11, §15.3].

Theorem 3.4. *The Hahn-Exton q -Bessel function of order $\nu > -1$ has a countably infinite number of positive simple zeros.*

Proof: In view of Corollary 3.2 and Lemma 3.3 it remains to show that $J_\nu(\cdot; q^2)$, with $\nu > -1$, has a countably infinite number of zeros. We prove that the assumption $J_\nu(\cdot; q^2)$, with $\nu > -1$, has a finite number of real positive zeros, leads to a contradiction.

Assume that $a > 0$ is the largest positive zero of $J_\nu(\cdot; q^2)$ and choose $x > 0$ satisfying

$$x^2 q^2 - 1 - q^{2\nu} > 0, \quad \text{and} \quad q^2 x > a.$$

Hence, $J_\nu(xq^2; q^2)$, $J_\nu(xq; q^2)$ and $J_\nu(x; q^2)$ are non-zero real numbers of the same sign and

$$J_\nu(xq^2; q^2) + q^{-\nu} (q^2 x^2 - 1 - q^{2\nu}) J_\nu(xq; q^2) + J_\nu(x; q^2)$$

is a non-zero real number. This is in contradiction with the second order q -difference equation (2.12) for the Hahn-Exton q -Bessel function. $\square$

Theorem 3.4 shows that we can order the positive zeros of $J_\nu(\cdot; q^2)$, with $\nu > -1$, as

$$0 < j_1^\nu(q^2) < j_2^\nu(q^2) < j_3^\nu(q^2) < \dots \quad (3.4)$$

The positive zeros of $J_\nu(\cdot; q^2)$ will also be denoted by j_n or j_n^ν . Proposition 3.1 and relation (3.3) can be combined to state the Fourier-Bessel orthogonality relations for the Hahn-Exton q -Bessel function, cf. [1] and [2].

Proposition 3.5. *Let $\nu > -1$ and $0 < j_1 < j_2 < \dots$ be the positive zeros of the Hahn-Exton q -Bessel function $J_\nu(\cdot; q^2)$, then*

$$\begin{aligned} & \int_0^1 x J_\nu(qj_n x; q^2) J_\nu(qj_m x; q^2) d_q x \\ &= -\frac{1}{2} (1-q) q^{\nu-1} J_{\nu+1}(qj_n; q^2) J'_\nu(j_n; q^2) \delta_{n,m} \\ &= \frac{1}{2} (1-q)^2 q^{\nu-2} \left(D_q J_\nu(\cdot; q^2) \right) (j_n) J'_\nu(j_n; q^2) \delta_{n,m} \\ &= -\frac{1}{2} (1-q) q^{\nu-2} j_n^{-1} J_\nu(qj_n; q^2) J'_\nu(j_n; q^2) \delta_{n,m} \\ &= -\frac{1}{2} (1-q) q^{-2} J_{\nu+1}(j_n; q^2) J'_\nu(j_n; q^2) \delta_{n,m}. \end{aligned}$$

Proof: It remains to prove the last three equalities. The first one is a consequence of

$$\left[D_q J_\nu(\cdot; q^2) \right] (x) = \frac{-q}{1-q} J_{\nu+1}(xq; q^2) + \frac{1-q^\nu}{x(1-q)} J_\nu(x; q^2), \quad (3.5)$$

which follows from (2.6) and (2.11). The second equality follows from the definition of the q -derivative D_q , cf. (2.5). The third equality follows from $J_{\nu+1}(j_n; q^2) = q^{\nu+1} J_{\nu+1}(qj_n; q^2)$, which is a consequence of (2.13) and (2.14). $\square$

Next we will derive that the zeros of the Hahn-Exton q -Bessel functions of order ν and $\nu+1$ are interlaced similarly to the interlacing property of the zeros of the Bessel functions of order ν and $\nu+1$, cf. Watson [11, §15.22]. We start with the following lemma.

Lemma 3.6. *Let $\nu > -1$, then $J_{\nu-1}(\cdot; q^2)$ and $J_{\nu+1}(\cdot; q^2)$ each have at least one zero between two consecutive positive zeros of $J_\nu(\cdot; q^2)$.*

Proof: Consider the function

$$g(x) = \frac{x^\nu}{1-q} J_{\nu-1}(x; q^2) = D_q \left((\cdot)^\nu J_\nu(\cdot; q^2) \right) (x),$$

where we used (2.10) as well. If a is a positive zero of $J_\nu(\cdot; q^2)$, then

$$g(a) = -\frac{(aq)^\nu J_\nu(aq; q^2)}{(1-q)a}$$

by (2.5). It follows from Proposition 3.5 and Lemma 3.3 that $g(a)g(b) < 0$ for two consecutive zeros $0 < a < b$ of $J_\nu(\cdot; q^2)$. This proves the lemma for $J_{\nu-1}(\cdot; q^2)$.

The other case is reduced to the result for $J_{\nu-1}(\cdot; q^2)$, since $J_{\nu-1}(a; q^2) = -J_{\nu+1}(a; q^2)$ for a non-zero zero of $J_\nu(\cdot; q^2)$ by (2.13). $\square$

Theorem 3.7. *The positive real zeros of $J_\nu(\cdot; q^2)$ and $J_{\nu+1}(\cdot; q^2)$, with $\nu > -1$, interlace. Explicitly,*

$$0 < j_1^\nu < j_1^{\nu+1} < j_2^\nu < j_2^{\nu+1} < j_3^\nu < j_3^{\nu+1} < \dots$$

Proof: The interlacing of the zeros of $J_\nu(\cdot; q^2)$ and $J_{\nu+1}(\cdot; q^2)$ is a straightforward application of Lemma 3.6. It remains to prove that $j_1^\nu < j_1^{\nu+1}$.

First note that the ${}_1\phi_1$ -series in (2.4) yields 1 for $x = 0$. Hence, by continuity,

$$J_\nu(x; q^2) > 0, \quad x \in (0, j_1^\nu), \quad \nu > -1. \quad (3.6)$$

Thus $J'_\nu(j_1^\nu; q^2) < 0$ and by Lemma 3.3 and Proposition 3.5 we get $J_{\nu+1}(j_1^\nu; q^2) > 0$. From (3.6) it follows that there exists an even number of positive zeros of $J_{\nu+1}(\cdot; q^2)$ in the open interval $(0, j_1^\nu)$. Lemma 3.6 and j_1^ν being the smallest positive zero of $J_\nu(\cdot; q^2)$ imply that this number is zero. $\square$

An upper bound for the first zero j_1^ν may be obtained as follows. Take $x = j_1^\nu$ in (2.12) to obtain

$$J_\nu(j_1^\nu q^2; q^2) + q^{-\nu} \left((j_1^\nu)^2 q^2 - 1 - q^{2\nu} \right) J_\nu(j_1^\nu q; q^2) = 0.$$

Inequality (3.6) implies that both q -Bessel functions are positive, so that

$$(j_1^\nu)^2 q^2 - 1 - q^{2\nu} < 0 \iff j_1^\nu < q^{-1} \sqrt{1 + q^{2\nu}}.$$

Theorem 3.7 shows that for $\nu > -1$ the Hahn-Exton q -Bessel functions of order ν and $\nu + 1$ do not have zeros in common except possibly 0. This also holds for general order.

Proposition 3.8. *The Hahn-Exton q -Bessel functions $J_\nu(\cdot; q^2)$ and $J_{\nu+1}(\cdot; q^2)$ have no common zeros except possibly 0.*

Proof: We argue by contradiction, so let $x \in \mathbb{C}$ be a non-zero zero of both $J_\nu(\cdot; q^2)$ and $J_{\nu+1}(\cdot; q^2)$. Eliminating $J_{\nu+1}(x; q^2)$ from (2.13) and (2.14) leads to

$$J_{\nu+1}(x; q^2) - q^{\nu+1} J_{\nu+1}(xq; q^2) = x J_\nu(x; q^2).$$

Hence, $J_{\nu+1}(xq; q^2) = 0$ as well. Next consider the q -difference equation (2.12) with ν replaced by $\nu + 1$. Since x and xq are zeros of $J_{\nu+1}(\cdot; q^2)$ we find $J_{\nu+1}(xq^2; q^2) = 0$. Repeated application of (2.12) leads to $J_{\nu+1}(xq^k; q^2) = 0$ for all $k \in \mathbb{Z}_+$. This implies $J_{\nu+1}(\cdot; q^2) \equiv 0$, which is the desired contradiction. $\square$

Finally we state a result on the zeros of the q -derivative of the Hahn-Exton q -Bessel function. Specialise $z = 1$ in Proposition 3.1 and use (3.5) to obtain

$$\begin{aligned} (a^2 - b^2) \int_0^1 x J_\nu(aqx; q^2) J_\nu(bqx; q^2) d_q x \\ = (1 - q)^2 q^{\nu-2} \left(b \left(D_q J_\nu(\cdot; q^2) \right) (b) J_\nu(a; q^2) - a \left(D_q J_\nu(\cdot; q^2) \right) (a) J_\nu(b; q^2) \right) \end{aligned} \quad (3.7)$$

for $\Re(\nu) > -1$ and $a, b \in \mathbb{C} \setminus \{0\}$.

Proposition 3.9. *The non-zero zeros of $D_q J_\nu(\cdot; q^2)$ for $\nu > 0$ are real and simple.*

Proof: In order to prove that the zeros of $D_q J_\nu(\cdot; q^2)$ are real, we proceed as in the proof of Corollary 3.2. In this case purely imaginary zeros can be excluded for $\nu > 0$.

To see that the zeros are simple, we apply l'Hôpital's rule to (3.7) and then take a a non-zero real zero of $D_q J_\nu(\cdot; q^2)$. The result is

$$\int_0^1 x \left| J_\nu(aqx; q^2) \right|^2 d_q x = -\frac{1}{2} (1 - q)^2 q^{\nu-2} J_\nu(a; q^2) \left(D_q J_\nu(\cdot; q^2) \right)'(a).$$

Since the left hand side is non-zero, the result follows. $\square$

4 On the associated q -Lommel polynomials.

The Bessel function $J_\nu(z)$ satisfies the three term recurrence relation

$$J_{\nu+1}(z) = \frac{2\nu}{z} J_\nu(z) - J_{\nu-1}(z). \quad (4.1)$$

If we iterate relation (4.1), we can express $J_{\nu+m}(z)$, with $m \in \mathbb{Z}_+$, in terms of $J_\nu(z)$ and $J_{\nu-1}(z)$, with coefficients that are polynomials in $\frac{1}{z}$. Indeed we have

$$J_{\nu+m}(z) = R_{m,\nu}(z)J_\nu(z) - R_{m-1,\nu+1}(z)J_{\nu-1}(z), \quad (4.2)$$

where $R_{m,\nu}(z)$ are the Lommel polynomials, see Watson [11, §9.6]. These polynomials satisfy the three term recurrence relation, cf. [11, 9.63(2)]

$$R_{m+1,\nu}(z) = \frac{2(\nu+m)}{z}R_{m,\nu}(z) - R_{m-1,\nu}(z), \quad (4.3)$$

with $R_{0,\nu}(z) = 1$ and $R_{1,\nu}(z) = \frac{2\nu}{z}$. Usually a related set of polynomials, the so-called modified Lommel polynomials, is defined by

$$h_{m,\nu}(z) = R_{m,\nu}\left(\frac{1}{z}\right).$$

It is clear that $h_{m,\nu}(z)$ is a polynomial in z of degree m , which satisfies the three term recurrence relation

$$h_{m+1,\nu}(z) = 2z(\nu+m)h_{m,\nu}(z) - h_{m-1,\nu}(z).$$

By Favard's theorem, the modified Lommel polynomials are orthogonal with respect to a positive measure. The orthogonality measure is a discrete measure with weights at $\frac{1}{j_n^\nu}$, where j_n^ν are the zeros of the Bessel function $J_\nu(z)$.

The asymptotic behaviour of the Lommel polynomials $R_{m,\nu}$ is related to the Bessel function by Hurwitz's formula, cf. [11, 9.65(1)]

$$\frac{\left(\frac{1}{2}z\right)^{\nu+m} R_{m,\nu+1}(z)}{\Gamma(\nu+m+1)} \xrightarrow{m \rightarrow \infty} J_\nu(z). \quad (4.4)$$

In the next subsections we will derive and investigate two different q -analogues of the Lommel polynomials. The first type follows from the recurrence relation (2.13). The second type is a result of the difference-recurrence relation (2.14).

4.1 q -Lommel polynomials associated with (2.13)

In this subsection we determine q -analogues of the Lommel polynomials arising from the recurrence relation (2.13). We will present an explicit formula, a generating function and an analogue of Hurwitz's formula (4.4). In order to derive q -analogues of the results mentioned in the previous section, we introduce a second Hahn-Exton q -Bessel function. Define

$$\begin{aligned} \mathcal{J}_\nu(x; q) &= e^{i\nu\pi} q^{-\frac{1}{2}\nu} J_{-\nu}(xq^{-\frac{1}{2}\nu}; q), \\ &= e^{i\nu\pi} \frac{(q^{-\nu+1}; q)_\infty}{(q; q)_\infty} x^{-\nu} q^{\frac{1}{2}\nu(\nu-1)} \sum_{k=0}^{\infty} \frac{(-1)^k q^{\frac{1}{2}k(k+1)} x^{2k} q^{-\nu k}}{(q^{-\nu+1}; q)_k (q; q)_k} \end{aligned} \quad (4.5)$$

Relations (2.10) and (2.11) give us, also in combination with the definition (4.5) above, the next formulas:

$$J_\nu(xq^{\frac{1}{2}}; q) = q^{\frac{1}{2}\nu} J_\nu(x; q) + xq^{\frac{1}{2}} J_{\nu+1}(xq^{\frac{1}{2}}; q), \quad (4.6)$$

$$\mathcal{J}_\nu(xq^{\frac{1}{2}}; q) = q^{\frac{1}{2}\nu} \mathcal{J}_\nu(x; q) + xq^{\frac{1}{2}} \mathcal{J}_{\nu+1}(xq^{\frac{1}{2}}; q), \quad (4.7)$$

$$J_\nu(xq^{\frac{1}{2}}; q) = q^{-\frac{1}{2}\nu} J_\nu(x; q) - xq^{-\frac{1}{2}\nu} J_{\nu-1}(x; q), \quad (4.8)$$

$$\mathcal{J}_\nu(xq^{\frac{1}{2}}; q) = q^{-\frac{1}{2}\nu} \mathcal{J}_\nu(x; q) - xq^{-\frac{1}{2}\nu} \mathcal{J}_{\nu-1}(x; q). \quad (4.9)$$

Furthermore, $\mathcal{J}_\nu(x; q^2)$ also satisfies the relations (2.12) and (2.13), i.e.

$$\left\{ x + \frac{1 - q^\nu}{x} \right\} \mathcal{J}_\nu(x; q) = \mathcal{J}_{\nu-1}(x; q) + \mathcal{J}_{\nu+1}(x; q), \quad (4.10)$$

$$\mathcal{J}_\nu(xq; q) + q^{-\frac{1}{2}\nu} (x^2 q - 1 - q^\nu) \mathcal{J}_\nu(xq^{\frac{1}{2}}; q) + \mathcal{J}_\nu(x; q) = 0. \quad (4.11)$$

This follows by combining relations (4.7) and (4.9).

Next, we will give two q -analogues of (4.2):

Proposition 4.1. *The functions $J_\nu(x; q)$ and $\mathcal{J}_\nu(x; q)$ satisfy the (same) recurrence relations*

$$J_{\nu+m}(x; q) = R_{m,\nu}(x; q) J_\nu(x; q) - R_{m-1,\nu+1}(x; q) J_{\nu-1}(x; q), \quad (4.12)$$

$$\mathcal{J}_{\nu+m}(x; q) = R_{m,\nu}(x; q) \mathcal{J}_\nu(x; q) - R_{m-1,\nu+1}(x; q) \mathcal{J}_{\nu-1}(x; q). \quad (4.13)$$

Proof: We will give the proof for $J_\nu(x; q)$. We can iterate the recurrence relation (2.13) to get an expression of the form

$$J_{\nu+m}(x; q) = R_{m,\nu}(x; q) J_\nu(x; q) - S_{m,\nu}(x; q) J_{\nu-1}(x; q).$$

Next consider $J_{\nu+m+1}(x; q)$. It satisfies the relations

$$\begin{aligned} J_{\nu+m+1}(x; q) &= R_{m+1,\nu}(x; q) J_\nu(x; q) - S_{m+1,\nu}(x; q) J_{\nu-1}(x; q) \\ &= R_{m,\nu+1}(x; q) J_{\nu+1}(x; q) - S_{m,\nu+1}(x; q) J_\nu(x; q) \\ &= \left[R_{m,\nu+1}(x; q) \left\{ x + \frac{1 - q^\nu}{x} \right\} - S_{m,\nu+1}(x; q) \right] J_\nu(x; q) - R_{m,\nu+1}(x; q) J_{\nu-1}(x; q). \end{aligned}$$

Hence $S_{m,\nu}(x; q) = R_{m-1,\nu+1}(x; q)$ and the proof is completed. Since $J_\nu(x; q)$ and $\mathcal{J}_\nu(x; q)$ satisfy the same recurrence relation (compare (2.13) and (4.10)), it is obvious that (4.13) will also hold. $\square$

Another consequence of the proof is the recurrence relation

$$R_{m+1,\nu}(x; q) = \left\{ x + \frac{1 - q^\nu}{x} \right\} R_{m,\nu+1}(x; q) - R_{m-1,\nu+2}(x; q). \quad (4.14)$$

Since (4.12) and (4.13) are q -analogues of (4.2), the functions $R_{m,\nu}$ are q -analogues of the Lommel polynomials. Later, when an explicit formula is derived, this will become clearer.

Lemma 4.2. *For non-integral ν the function $R_{m,\nu}$ can be expressed as*

$$R_{m,\nu}(x; q) = \frac{xe^{-i\pi\nu} q^{-\frac{1}{2}\nu(\nu-1)} (q; q)_\infty (q; q)_\infty}{(q^\nu; q)_\infty (q^{-\nu+1}; q)_\infty} \times \{J_{\nu-1}(x; q) J_{\nu+m}(x; q) - J_{\nu+m}(x; q) J_{\nu-1}(x; q)\}. \quad (4.15)$$

Proof. Multiply the recurrence relations (4.12) and (4.13) with respectively $J_{\nu-1}(x; q)$ and $J_{\nu-1}(x; q)$ and subtract the resulting formulas to obtain

$$R_{m,\nu}(x; q) = \left\{ \frac{J_{\nu+m}(x; q) J_{\nu-1}(x; q) - J_{\nu+m}(x; q) J_{\nu-1}(x; q)}{J_{\nu-1}(x; q) J_\nu(x; q) - J_\nu(x; q) J_{\nu-1}(x; q)} \right\}.$$

Using (4.8) and (4.9) we rewrite the denominator of this expression as

$$\begin{aligned} & J_{\nu-1}(x; q) J_\nu(x; q) - J_\nu(x; q) J_{\nu-1}(x; q) \\ &= J_\nu(x; q) \left[\frac{1}{x} J_\nu(x; q) - \frac{1}{x} q^{\frac{1}{2}\nu} J_\nu(xq^{\frac{1}{2}}; q) \right] - J_\nu(x; q) \left[\frac{1}{x} J_\nu(x; q) - \frac{1}{x} q^{\frac{1}{2}\nu} J_\nu(xq^{\frac{1}{2}}; q) \right] \\ &= \frac{q^{\frac{1}{2}\nu}}{x} \left\{ J_\nu(x; q) J_\nu(xq^{\frac{1}{2}}; q) - J_\nu(x; q) J_\nu(xq^{\frac{1}{2}}; q) \right\}. \end{aligned}$$

The term in braces is a Wronskian type formula, which has been evaluated by R.F. Swarttouw [9, (4.3.8)]. Explicitly,

$$J_\nu(x; q) J_\nu(xq^{\frac{1}{2}}; q) - J_\nu(x; q) J_\nu(xq^{\frac{1}{2}}; q) = q^{\frac{1}{2}\nu(\nu-2)} e^{i\nu\pi} \frac{(q^\nu; q)_\infty (q^{-\nu+1}; q)_\infty}{(q; q)_\infty (q; q)_\infty}, \quad (4.16)$$

which is non-zero for non-integral values of ν . Now the lemma follows easily. $\square$

Remark: The case $\nu = n \in \mathbb{Z}$ requires a closer investigation, since (4.16) equals zero if $\nu \in \mathbb{Z}$. However, for $n \in \mathbb{Z}$ we have the relation, cf. [9, (3.2.12)]

$$J_n(x; q) = (-1)^n q^{-\frac{1}{2}n} J_{-n}(xq^{-\frac{1}{2}n}; q) = J_n(x; q), \quad (4.17)$$

so that the numerator in (4.15) will also become zero for $n \in \mathbb{Z}$. Now we can apply l'Hôpital's rule to (4.15) to show that the formula still makes sense for $\nu \in \mathbb{Z}$. However, it will be easier to show that $R_{m,n}(x; q)$ is the analytic continuation of $R_{m,\nu}(x; q)$, when we have the explicit representation for the $R_{m,\nu}(x; q)$. Until (4.23) we assume that ν is non-integral, but it is easily verified by analytic continuation that the results obtained in between are valid for integer ν as well.

Relation (4.15) gives rise to a three term recurrence relation for the q -Lommel polynomials:

Proposition 4.3. *The function $R_{m,\nu}(x; q)$ satisfies the recurrence relation*

$$\left\{x^2 + 1 - q^{\nu+m}\right\} R_{m,\nu}(x; q) = x \{R_{m-1,\nu}(x; q) + R_{m+1,\nu}(x; q)\}. \quad (4.18)$$

with $R_{0,\nu}(x; q) = 1$ and $R_{1,\nu}(x; q) = x + \frac{1-q^\nu}{x}$.

Proof: $R_{0,\nu}(x; q)$ and $R_{1,\nu}(x; q)$ follow immediately from (4.13) and (4.10). In order to shorten the notation we introduce

$$C_\nu(x; q) = \frac{x e^{-i\nu\pi} q^{-\frac{1}{2}\nu(\nu-1)}(q; q)_\infty(q; q)_\infty}{(q^\nu; q)_\infty(q^{-\nu+1}; q)_\infty}. \quad (4.19)$$

Now, starting with the lefthand side of (4.15), with m replaced by $m+1$, we find with the recurrence relations (2.13) and (4.10), with ν replaced by $\nu+m$

$$\begin{aligned} R_{m+1,\nu}(x; q) &= C_\nu(x; q) \{J_{\nu-1}(x; q) \mathcal{J}_{\nu+m+1}(x; q) - J_{\nu+m+1}(x; q) \mathcal{J}_{\nu-1}(x; q)\} \\ &= C_\nu(x; q) \left\{ J_{\nu-1}(x; q) \left[\left\{ x + \frac{1-q^{\nu+m}}{x} \right\} \mathcal{J}_{\nu+m}(x; q) - \mathcal{J}_{\nu+m-1}(x; q) \right] + \right. \\ &\quad \left. - \mathcal{J}_{\nu-1}(x; q) \left[\left\{ x + \frac{1-q^{\nu+m}}{x} \right\} J_{\nu+m}(x; q) - J_{\nu+m-1}(x; q) \right] \right\} \\ &= \left\{ x + \frac{1-q^{\nu+m}}{x} \right\} R_{m,\nu}(x; q) - R_{m-1,\nu}(x; q) \end{aligned}$$

which proves the proposition. $\square$

From Proposition 4.3 it is easy to see that the function $R_{m,\nu}$ is not a polynomial, but a function of the form

$$R_{m,\nu}(x; q) = \sum_{n=-m}^m c_n x^n.$$

So $R_{m,\nu}$ is in fact a Laurent polynomial. This suggests the introduction of the function $p_m(x; q) = x^m R_{m,\nu}(x; q)$. From (4.18) it follows that $p_m(x; q)$ satisfies the three term recurrence relation

$$\begin{aligned} (x^2 + 1 - q^{\nu+m}) p_m(x; q) &= p_{m+1}(x; q) + x^2 p_{m-1}(x; q), \\ p_0(x; q) &= 1 \quad \text{and} \quad p_1(x; q) = x^2 + (1 - q^\nu). \end{aligned} \quad (4.20)$$

It follows from (4.20) that p_m is actually a polynomial of degree m in x^2 . The polynomial p_m does not satisfy the conditions of Favard's theorem, cf. [10, II.3.2], so that it is not orthogonal with respect to a positive measure. However, $p_m(x; q)$ possesses nice properties, which are q -analogues of the well-known properties of the Lommel polynomials.

First we will derive a generating function for the q -Lommel polynomials. Multiply (4.20) with t^{m+1} and sum from $m=1$ to ∞ . When we introduce the generating function

$$G(x, t) = \sum_{m=0}^{\infty} t^m p_m(x; q),$$

we find

$$\begin{aligned} & x^2 t (G(x, t) - 1) + t (G(x, t) - 1) - q^\nu t (G(x, qt) - 1) \\ &= G(x, t) - 1 - t (x^2 + 1 - q^\nu) + x^2 t^2 G(x, t), \end{aligned}$$

which simplifies to

$$G(x, t) = \frac{1 - q^\nu t G(x, qt)}{(1 - t)(1 - x^2 t)}. \quad (4.21)$$

When we iterate relation (4.21) and make use of the fact that $G(x, 0) = 1$, we have

$$G(x, t) = \sum_{j=0}^{\infty} \frac{q^{j\nu} q^{\frac{1}{2}j(j-1)} (-t)^j}{(t; q)_{j+1} (x^2 t; q)_{j+1}}.$$

This expression is valid for $t \neq q^{-p}$, $x^2 t \neq q^{-p}$, $p \in \mathbb{Z}_+$. So we have the following generating function for the function $R_{m,\nu}(x; q)$:

$$\begin{aligned} \sum_{m=0}^{\infty} (xt)^m R_{m,\nu}(x; q) &= \sum_{j=0}^{\infty} \frac{q^{j\nu} q^{\frac{1}{2}j(j-1)} (-t)^j}{(t; q)_{j+1} (tx^2; q)_{j+1}} \\ &= \frac{1}{(1 - t)(1 - x^2 t)^{2\phi_2}} \left(\begin{matrix} q, 0 \\ qt, qtx^2 \end{matrix} \middle| q; tq^\nu \right). \end{aligned} \quad (4.22)$$

The generating function (4.22) gives rise to an explicit expression of $R_{m,\nu}(x; q)$. Use the q -binomial theorem [3, (1.3.2)] twice to obtain

$$\begin{aligned} \frac{1}{(t; q)_{j+1}} &= \frac{(tq^{j+1}; q)_\infty}{(t; q)_\infty} = {}_1\phi_0 \left(\begin{matrix} tq^{j+1} \\ - \end{matrix} \middle| q; t \right), \quad |t| < 1, \\ \frac{1}{(tx^2; q)_{j+1}} &= {}_1\phi_0 \left(\begin{matrix} q^{j+1} \\ - \end{matrix} \middle| q; tx^2 \right), \quad |tx^2| < 1. \end{aligned}$$

This yields

$$\sum_{m=0}^{\infty} (xt)^m R_{m,\nu}(x; q) = \sum_{j,k,n=0}^{\infty} \frac{q^{j\nu} q^{\frac{1}{2}j(j-1)} (-1)^j (q^{j+1}; q)_k (q^{j+1}; q)_n}{(q; q)_k (q; q)_n} x^{2n} t^{j+k+n}.$$

Equating powers of t yields

$$\begin{aligned} x^m R_{m,\nu}(x; q) &= \sum_{n=0}^m \frac{x^{2n}}{(q; q)_n} \sum_{j=0}^{m-n} \frac{q^{j\nu} q^{\frac{1}{2}j(j-1)} (-1)^j (q^{j+1}; q)_{m-n-j} (q^{j+1}; q)_n}{(q; q)_{m-n-j}} \\ &= \sum_{n=0}^m x^{2n} \sum_{j=0}^{m-n} \frac{(q^{n-m}; q)_j (q^{n+1}; q)_j}{(q; q)_j (q; q)_j} q^{j\nu} q^{j(m-n)} \\ &= \sum_{n=0}^m x^{2n} {}_2\phi_1 \left(\begin{matrix} q^{n-m}, q^{n+1} \\ q \end{matrix} \middle| q; q^{\nu+m-n} \right). \end{aligned}$$

This expression for $R_{m,\nu}(x; q)$ shows the analytic dependence on q^ν , cf. the remark following Lemma 4.2. Finally, using Heine's transformation formula, cf. [3, (1.4.1)], we obtain the explicit representation

$$R_{m,\nu}(x; q) = \sum_{n=0}^m x^{2n-m} \frac{(q^{n+1}; q)_\infty (q^\nu; q)_\infty}{(q; q)_\infty (q^{\nu+m-n}; q)_\infty} {}_2\phi_1 \left(\begin{matrix} q^{-n}, q^{\nu+m-n} \\ q^\nu \end{matrix} \middle| q; q^{n+1} \right). \quad (4.23)$$

The explicit expression (4.23) can also be obtained from (4.15) in combination with the explicit series representations (4.5) and (2.4). In this case a formula [9, (6.4.4)] for the product of two Hahn-Exton q -Bessel functions has to be used.

Formally we can obtain a q -analogue of Hurwitz's formula (4.4) by taking termwise limits in the explicit representation (4.23). This gives

$$\begin{aligned} x^m R_{m,\nu}(x; q) &\xrightarrow{m \rightarrow \infty} \frac{(q^\nu; q)_\infty}{(q; q)_\infty} \sum_{n=0}^{\infty} x^{2n} (q^{n+1}; q)_\infty {}_2\phi_1 \left(\begin{matrix} q^{-n}, 0 \\ q^\nu \end{matrix} \middle| q; q^{n+1} \right) \\ &= (q^{\nu+1}; q)_\infty \sum_{k=0}^{\infty} \frac{q^k}{(q^\nu; q)_k (q; q)_k} \sum_{n=k}^{\infty} \frac{(q^{-n}; q)_k}{(q; q)_n} q^{nk} x^{2n}. \end{aligned}$$

The inner sum equals

$$(-1)^k q^{\frac{1}{2}k(k-1)} x^{2k} \frac{1}{(x^2; q)_\infty},$$

so that we formally obtain

$$\begin{aligned} x^m R_{m,\nu}(x; q) &\xrightarrow{m \rightarrow \infty} \frac{(q^\nu; q)_\infty}{(x^2; q)_\infty} {}_1\phi_1 \left(\begin{matrix} 0 \\ q^\nu \end{matrix} \middle| q; qx^2 \right) \\ &= \frac{(q; q)_\infty}{(x^2; q)_\infty} x^{1-\nu} J_{\nu-1}(x; q). \end{aligned} \quad (4.24)$$

4.2 q -Lommel polynomials associated with (2.14)

In this subsection we consider the q -analogues of the Lommel polynomials that arise from the iteration of (2.14). Note that the shift in the argument in (2.14) will present some difficulty. As a consequence, there is no unique definition of these q -Lommel polynomials. For a certain choice we show that the (modified) q -Lommel polynomials associated with the Jackson q -Bessel function, see M.E.H. Ismail [4], are obtained in this context. These polynomials are orthogonal as shown by M.E.H. Ismail [4, (4.16)]. For a slightly different definition of the q -Lommel polynomials we present a Hurwitz-type formula, cf. (4.4).

We start with iterating (2.14).

Lemma 4.4. *There exist unique constants $a_i(\nu, m)$ and $b_j(\nu, m)$ so that*

$$J_{\nu+m}(xq^m; q^2) = \sum_{i=0}^{[m/2]} \frac{a_i(\nu, m)}{x^{m-2i}} J_\nu(xq^i; q^2) + \sum_{j=0}^{[(m-1)/2]} \frac{b_j(\nu, m)}{x^{m-1-2j}} J_{\nu-1}(xq^j; q^2),$$

where $[a]$ denotes the largest integer smaller than or equal to a .

Proof: The proof uses induction with respect to m . The case $m = 0$ is trivial and the case $m = 1$ is just (2.14). For the induction step we use, cf. (2.14),

$$\begin{aligned} J_{\nu+m+1}(xq^{m+1}; q^2) &= J_{(\nu+m)+1}(xq^m \cdot q; q^2) \\ &= q^{-(\nu+m+1)} \frac{1 - q^{2(\nu+m)}}{xq^m} J_{\nu+m}(xq^m; q^2) - q^{-(\nu+m+1)} J_{\nu+(m-1)}(xq \cdot q^{m-1}; q^2). \end{aligned} \quad (4.25)$$

Use the induction hypothesis in the last two terms of (4.25) and shift summation parameters to prove the lemma. $\square$

Actually this proof yields a recurrence relation for the constants $a_i(\nu, m+1)$. Explicitly,

$$a_i(\nu, m+1) = q^{-(\nu+1)} q^{-2m} (1 - q^{2(\nu+m)}) a_i(\nu, m) - q^{-\nu+2(i-1-m)} a_{i-1}(\nu, m-1), \quad (4.26)$$

with $a_i(\nu, m) = 0$ for $i < 0$ or $i > [m/2]$. A similar result holds for the constants $b_j(\nu, m)$, but these constants will be related to the $a_i(\nu, m)$ in due course, cf. Lemma 4.5. So we do not present the recurrence relation for the $b_j(\nu, m)$'s.

Although Lemma 4.4 also holds with J_ν replaced by $\mathcal{J}_\nu$, cf. (4.5), this does not lead to nice expressions as in the previous subsection due to the shift in the argument in Lemma 4.4.

Motivated by Lemma 4.4 and the recurrence relation (4.26) we define

$$r_{m,\nu}(x; q^2) = \sum_{i=0}^{[m/2]} \frac{a_i(\nu, m)}{x^{m-2i}} q^{-i(i+1)}, \quad (4.27)$$

which is a polynomial of degree m in $\frac{1}{x}$. The factor $q^{-i(i+1)}$ in (4.27) is chosen in such a way that (4.26) yields a three term recurrence relation for the $r_{m,\nu}(x; q^2)$. Explicitly

$$q^{2m} r_{m+1,\nu}(x; q^2) = q^{-(\nu+1)} \frac{1 - q^{2(\nu+m)}}{x} r_{m,\nu}(x; q^2) - q^{-(\nu+2)} r_{m-1,\nu}(x; q^2), \quad (4.28)$$

which follows from multiplying (4.26) by $x^{-m-1+2i} q^{-i(i+1)}$ and summing from $i = 0$ to $[(m+1)/2]$. The begin values of the recurrence relation (4.28) are

$$r_{0,\nu}(x; q^2) = 1, \quad r_{1,\nu}(x; q^2) = q^{-(\nu+1)} (1 - q^{2\nu}) x^{-1}. \quad (4.29)$$

In order to determine the constants $a_i(\nu, m)$ in (4.27) we introduce the polynomial $h_{m,\nu}(x; q^2) = r_{m,\nu}(\frac{1}{x}; q^2)$. The three term recurrence relation (4.26) with its begin values (4.29) is rewritten as

$$\begin{aligned} q^{2m} h_{m+1,\nu}(x; q^2) &= q^{-\nu-1} (1 - q^{2(\nu+m)}) x h_{m,\nu}(x; q^2) - q^{-\nu-2} h_{m-1,\nu}(x; q^2), \\ h_{0,\nu}(x; q^2) &= 1 \quad \text{and} \quad h_{1,\nu}(x; q^2) = q^{-\nu-1} (1 - q^{2\nu}) x. \end{aligned} \quad (4.30)$$

Let us recall the (modified) q -Lommel polynomials associated with the Jackson q -Bessel function as presented by M.E.H. Ismail [4]. Define, cf. [4, (3.6)],

$$\tilde{h}_{m,\nu}(x; q^2) = \sum_{j=0}^{[m/2]} \frac{(2x)^{m-2j} (-1)^j (q^{2\nu}; q^2)_{m-j} (q^2; q^2)_{m-j}}{(q^2; q^2)_j (q^{2\nu}; q^2)_j (q^2; q^2)_{m-2j}} q^{2j(j+\nu-1)}, \quad (4.31)$$

then the set of polynomials $\tilde{h}_{m,\nu}(x; q^2)$ satisfies the three term recurrence relation, cf. [4, (1.22)],

$$\begin{aligned} \tilde{h}_{m+1,\nu}(x; q^2) &= 2x \left(1 - q^{2(\nu+m)}\right) \tilde{h}_{m,\nu}(x; q^2) - q^{2(m+\nu-1)} \tilde{h}_{m-1,\nu}(x; q^2), \\ \tilde{h}_{0,\nu}(x; q^2) &= 1 \quad \text{and} \quad \tilde{h}_{1,\nu}(x; q^2) = 2x \left(1 - q^{2\nu}\right). \end{aligned} \quad (4.32)$$

A straightforward calculation shows that (4.30) and (4.32) are equivalent. It suffices to take

$$h_{m,\nu}(x; q^2) = q^{-\frac{3}{2}m\nu - m^2} \tilde{h}_{m,\nu}(q^{\frac{\nu}{2}} x/2; q^2). \quad (4.33)$$

From (4.33), (4.31) and (4.27) we obtain the explicit expression

$$a_i(m, \nu) = q^{-m(m+\nu)} q^{i(3i+\nu-1)} \frac{(-1)^i (q^{2\nu}; q^2)_{m-i} (q^2; q^2)_{m-i}}{(q^2; q^2)_i (q^{2\nu}; q^2)_i (q^2; q^2)_{m-2i}}. \quad (4.34)$$

Now that we have determined the value of $a_i(\nu, m)$ we present its link to the constants $b_j(\nu, m)$.

Lemma 4.5.

$$b_j(\nu, m+1) = -q^{2j-m-\nu-1} a_j(\nu+1, m). \quad (4.35)$$

Proof: Consider

$$J_{\nu+m+1}(xq^{m+1}; q^2) = J_{(\nu+1)+m}(xq \cdot q^m; q^2) \quad (4.36)$$

and apply Lemma 4.4 with ν replaced by $\nu+1$ to the right hand side of (4.36). In the resulting sum involving the Hahn-Exton q -Bessel function of order $\nu+1$ we use (2.14). Comparison of the coefficients of $x^{2i-m} J_{\nu-1}(xq^i; q^2)$ with the ones obtained by applying Lemma 4.4 with m replaced by $m+1$ to the left hand side of (4.36) proves the lemma. $\square$

In the proof of Lemma 4.5 we can also compare the coefficients of $x^{2i-m-1} J_{\nu}(xq^i; q^2)$ and this results in

$$a_i(\nu, m+1) = q^{i-m-\nu-1} \left(1 - q^{2\nu}\right) a_i(\nu+1, m) + q^{2i-m-1} b_{i-1}(\nu+1, m). \quad (4.37)$$

Using Lemma 4.5 and (4.27) we can rewrite (4.37) as

$$\begin{aligned} r_{m+1,\nu}(x; q^2) &= \frac{1}{x} \left(1 - q^{2\nu}\right) q^{-(\frac{1}{2}m+\nu+1)} r_{m,\nu+1}(xq^{\frac{1}{2}}; q^2) \\ &\quad - q^{-(m+\nu+3)} r_{m-1,\nu+2}(xq; q^2). \end{aligned} \quad (4.38)$$

Since we have identified $r_{m,\nu}(x; q^2)$ with q -analogues of Lommel polynomials associated with Jackson's q -Bessel functions, Hurwitz's formula, cf. M.E.H. Ismail [4, (3.10)], holds for $r_{m,\nu}(x; q^2)$. We want to have a q -analogue of Hurwitz's formula involving the Hahn-Exton q -Bessel function and in order to do so we make use of the following q -analogue of the Lommel polynomial:

$$\tilde{r}_{m,\nu}(x; q) = \sum_{i=0}^{[m/2]} \frac{a_i(m, \nu)}{x^{m-2i}} q^{-2i(i-1)}. \quad (4.39)$$

From (4.26) we see that it satisfies the recurrence relation

$$\tilde{r}_{m+1,\nu}(x; q^2) = x^{-1} q^{-\nu-1-2m} (1 - q^{2(\nu+m)}) \tilde{r}_{m,\nu}(x; q^2) - q^{1-\nu-3m} \tilde{r}_{m-1,\nu}(q^{-1}x; q^2). \quad (4.40)$$

It follows from Favard's theorem, cf. [10, II.3.2], that $\tilde{r}_{m,\nu}(\frac{1}{x}; q^2)$ does not yield a set of orthogonal polynomials. However, from (4.34) and (4.39) we get

$$\begin{aligned} \tilde{r}_{m,\nu}(x; q^2) &= x^{-m} q^{-m(m+\nu)} (q^{2\nu}; q^2)_m \\ &\times {}_5\phi_3 \left(\begin{matrix} q^{1-m}, -q^{1-m}, q^{-m}, -q^{-m}, 0 \\ q^{2\nu}, q^{2(1-m-\nu)}, q^{-2m} \end{matrix} \middle| q^2; x^2 q^{2-\nu} \right) \end{aligned} \quad (4.41)$$

and from (4.41) we obtain the following q -analogue of Hurwitz's formula

$$x^m q^{m(m+\nu)} \tilde{r}_{m,\nu}(x; q^2) \xrightarrow{m \rightarrow \infty} (q^2; q^2)_\infty x^{1-\nu} q^{\frac{1}{2}\nu(1-\nu)} J_{\nu-1}(xq^{\frac{\nu}{2}}; q^2),$$

by taking formal termwise limits in (4.41).

References

- [1] H. EXTON, A basic analogue of the Bessel-Clifford equation, *Jñānābha* **8** (1978), 49 - 56.
- [2] H. EXTON, “ q -Hypergeometric functions and applications.” Ellis Horwood, Chichester, 1983.
- [3] G. GASPER AND M. RAHMAN, “Basic Hypergeometric Series,” *Encyclopedia of Mathematics and its Applications* **35**, Cambridge University Press, Cambridge, 1990.
- [4] M.E.H. ISMAIL, The zeros of basic Bessel functions, the functions $J_{\nu+ax}(x)$, and associated orthogonal polynomials, *Journal of Mathematical Analysis and Applications* **86** (1982), 1-19.
- [5] H.T. KOELINK, On q -Bessel functions related to the quantum group of plane motions, Report W 91-26, University of Leiden, 1991.
- [6] T.H. KOORNWINDER AND R.F. SWARTTOUW, On q -analogues of the Hankel- and Fourier Transforms, *Transactions of the American Mathematical Society* **333** (1992), 445-461.

- [7] M. RAHMAN, A note on the orthogonality of Jackson's q -Bessel function, *Canadian Mathematical Bulletin* **32** (1989), 369-376.
- [8] I.N. SNEDDON, "Fourier transforms," McGraw-Hill, New York, 1951.
- [9] R.F. SWARTTOUW, The Hahn-Exton q -Bessel function, Ph. D. thesis, Delft Technical University, 1992.
- [10] G. SZEGÖ, "Orthogonal Polynomials," Colloquium publications 23, American Mathematical Society, 4th edition, Providence, Rhode Island, 1975.
- [11] G.N. WATSON, "A Treatise on the Theory of Bessel Functions," second edition, Cambridge University Press, Cambridge, 1966.